

**PHARMACOLOGICAL PROPERTIES OF ANTI-INFLAMMATORY AND
ANTIALLERGIC AGENTS: ANALYSIS FROM THE PERSPECTIVE OF
MICROBIOLOGICAL PROCESSES**

Komilova N.U.

Djanayev G.Yu.

Tashkent State Medical University, Tashkent, Uzbekistan

DOI: <https://doi.org/10.5281/zenodo.22915379>

Abstract. Inflammation and allergic reactions are important protective mechanisms of the human body, but excessive or prolonged activation of these processes can lead to tissue damage and various pathological conditions. Anti-inflammatory and antiallergic drugs play an important role in controlling inflammatory mediators, immune responses, and clinical manifestations of these disorders. From a microbiological perspective, inflammation is closely associated with interactions between microorganisms and the host immune system. Bacterial, viral, fungal, and other microbial agents can activate innate and adaptive immune responses through pathogen-associated molecular patterns and inflammatory mediators. The aim of this thesis is to analyze the pharmacological properties and mechanisms of action of anti-inflammatory and antiallergic agents in relation to microbiological and immunological processes.

Keywords: inflammation, allergy, anti-inflammatory drugs, antihistamines, glucocorticoids, NSAIDs, microorganisms, immune response, histamine, cytokines.

Introduction

Inflammation is a complex biological response to infection, tissue injury, or other harmful stimuli. Microorganisms and their structural components can activate immune cells and stimulate the release of inflammatory mediators such as cytokines, prostaglandins, leukotrienes, and histamine. Although inflammation has a protective role in eliminating infectious agents, excessive activation may contribute to tissue injury and chronic disease.

Allergic reactions represent another form of immune-mediated response in which the immune system reacts excessively to otherwise harmless environmental antigens. Mast cells, basophils, immunoglobulin E (IgE), histamine, leukotrienes, and other mediators are involved in the development of many allergic manifestations. Pharmacological intervention aims to reduce inflammation, inhibit mediator release, or block the effects of specific mediators.

Anti-inflammatory Agents and Their Mechanisms

Anti-inflammatory drugs can be broadly divided into non-steroidal anti-inflammatory drugs (NSAIDs) and glucocorticoids.

NSAIDs reduce inflammation primarily by inhibiting cyclooxygenase enzymes and decreasing the synthesis of prostaglandins. Reduced prostaglandin production contributes to decreased pain, fever, and inflammatory responses. Common representatives include ibuprofen, diclofenac, and naproxen. From a microbiological perspective, these agents do not directly eliminate bacteria, viruses, or fungi; rather, they modify the inflammatory response of the host. Therefore, in infectious diseases, symptomatic

Multidisciplinary research and Innovation (SICMRI 2026)

suppression of inflammation should not be considered a substitute for appropriate antimicrobial therapy when such therapy is indicated.

Glucocorticoids have a broader anti-inflammatory and immunosuppressive effect. They regulate gene transcription and reduce the production of several inflammatory cytokines and mediators. These drugs can decrease leukocyte activation, vascular permeability, and inflammatory tissue damage. However, prolonged or inappropriate use may suppress immune defense and increase susceptibility to certain infections. Therefore, their clinical use requires careful consideration of the underlying infectious and immunological status of the patient.

Antiallergic Agents

Antiallergic drugs are primarily used to control immune-mediated allergic reactions. Antihistamines are among the most commonly used agents. They primarily block H1 histamine receptors and reduce symptoms such as itching, sneezing, rhinorrhea, urticaria, and conjunctival irritation. Second-generation antihistamines, including cetirizine and loratadine, generally produce less central nervous system sedation than many first-generation agents.

Other antiallergic therapies include glucocorticoids, leukotriene receptor antagonists, and mast-cell stabilizing agents. These drugs act at different stages of the allergic inflammatory pathway and may be used according to the type and severity of the allergic condition.

Microbiological Perspective

The relationship between microorganisms and inflammatory responses is particularly important in clinical pharmacology. Microbial components can activate pattern-recognition receptors on immune and epithelial cells, leading to the release of cytokines and other inflammatory mediators. This response contributes to the elimination of pathogens but may also produce tissue damage when excessive.

The use of anti-inflammatory and antiallergic drugs can modify these host responses without directly affecting the causative microorganism. This distinction is clinically important. For example, NSAIDs may reduce fever and pain during infection, while glucocorticoids can significantly suppress inflammation and immune activity. Therefore, the potential influence of immunosuppressive therapy on host resistance to infection should be considered.

The interaction between the microbiota and the immune system is also relevant. Changes in the composition of normal microbial communities may influence immune regulation and inflammatory processes. Consequently, understanding microbiological and immunological mechanisms can contribute to more rational pharmacotherapy.

Conclusion

Anti-inflammatory and antiallergic agents are important components of modern pharmacotherapy. NSAIDs mainly reduce prostaglandin synthesis, while glucocorticoids suppress multiple inflammatory pathways. Antihistamines primarily inhibit the effects of histamine and reduce the clinical manifestations of allergic reactions.

From a microbiological perspective, these drugs primarily influence the host inflammatory and immune response rather than directly eliminating microorganisms. Therefore, their use in infectious and allergic conditions should be based on an understanding of the underlying pathophysiological and microbiological processes. Rational selection of therapy, appropriate dosing, and consideration of the patient's immune status are essential for achieving therapeutic effectiveness while minimizing potential complications.