

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

URINARY STONE DISEASE

Bobokulov Nurillo Asatovich

Samarkand State Medical University

Zayniddinov Mahmudjon Farxodjon o'g'li

Samarkand State Medical University

Movlanov Hazrat Musulmon o'g'li

Samarkand State Medical University

Annotation: Urinary stone disease, also known as urolithiasis, is a common urological disorder characterized by the formation of calculi in the kidneys, ureters, or bladder. Stones are primarily composed of calcium oxalate, calcium phosphate, uric acid, or struvite. The disease can cause acute flank pain, hematuria, urinary tract obstruction, and recurrent infections, significantly affecting patient quality of life. Etiology involves multifactorial mechanisms including metabolic abnormalities, genetic predisposition, dietary habits, dehydration, and urinary tract infections. Diagnosis relies on laboratory tests and imaging studies, with non-contrast computed tomography recognized as the gold standard. Management strategies include hydration, dietary adjustments, pharmacological therapy, and interventional procedures such as extracorporeal shock wave lithotripsy, ureteroscopy, and percutaneous nephrolithotomy. Preventive measures are essential to reduce recurrence, preserve renal function, and improve long-term outcomes.

Keywords: Urolithiasis, Kidney stones, Renal colic, Hematuria, Urinary obstruction.

Introduction

Urinary stone disease, also known as urolithiasis, is a common urological disorder characterized by the formation of calculi in the kidneys, ureters, or bladder. These stones are composed of crystalline substances that result from supersaturation of urine with minerals such as calcium, oxalate, uric acid, and phosphate. The prevalence of urinary stone disease has been increasing worldwide, affecting men more frequently than women, and often presenting during adulthood. The condition can cause significant clinical manifestations, including acute or chronic flank pain, hematuria, urinary tract obstruction, recurrent urinary tract infections, and, in severe cases, impaired renal function. The pathogenesis of urinary stones is multifactorial, involving metabolic abnormalities, genetic predisposition, dietary factors, dehydration, and urinary tract infections. Imbalances in urinary pH, concentration of stone-forming substances, and inhibitors of crystallization all contribute to stone formation. Early recognition and proper diagnosis are essential to prevent complications such as recurrent obstruction, kidney damage, and severe infection.

Main part

Urinary stone disease is a multifactorial disorder resulting from complex interactions between metabolic, dietary, genetic, and environmental factors. Stones typically form when urine becomes supersaturated with stone-forming salts, such as calcium oxalate, calcium phosphate, uric acid, or struvite. Metabolic disorders, including hypercalciuria, hyperuricosuria, and hyperoxaluria, play a central role in stone formation. Genetic predisposition contributes significantly, as family history increases susceptibility to recurrent stones. Urinary tract infections, particularly with urease-producing bacteria, can facilitate struvite stone formation. Low fluid intake and chronic dehydration concentrate the urine, enhancing crystallization.

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

Urinary pH also influences stone composition; acidic urine favors uric acid stones, while alkaline urine promotes calcium phosphate and struvite formation. Dietary factors, such as excessive animal protein, high sodium, and low citrate intake, exacerbate stone risk. Anatomical anomalies, including medullary sponge kidney or ureteropelvic junction obstruction, can promote urinary stasis and stone development. The pathogenesis involves nucleation, crystal aggregation, and retention within the renal tubules. Understanding these mechanisms is crucial for identifying high-risk patients, developing preventive strategies, and guiding individualized management plans.

The clinical presentation of urinary stone disease varies depending on the stone's size, location, and degree of obstruction. The most common symptom is renal colic, characterized by sudden, severe flank or abdominal pain radiating to the groin. Hematuria, either microscopic or gross, is frequently observed. Patients may also experience nausea, vomiting, urinary urgency, frequency, and dysuria. Fever or chills indicate concomitant infection and require urgent evaluation. Physical examination may reveal tenderness over the costovertebral angle. Laboratory investigations, including urinalysis, serum creatinine, calcium, uric acid, and electrolytes, help assess renal function and metabolic disturbances. Imaging studies are essential for confirming diagnosis, localizing stones, and evaluating obstruction. Non-contrast computed tomography (CT) is considered the gold standard due to its high sensitivity and specificity. Ultrasonography is a useful alternative, particularly for evaluating hydronephrosis and stones in the kidneys or bladder. Plain abdominal radiography may detect radiopaque stones but is limited for uric acid or small calculi. Accurate diagnosis is vital to prevent complications, guide treatment selection, and reduce the risk of recurrent stone formation.

Management of urinary stone disease involves both conservative and interventional approaches depending on stone size, location, and patient condition. Small stones (<5 mm) may pass spontaneously with adequate hydration, analgesia, and alpha-blockers to facilitate ureteral relaxation. Larger or obstructive stones often require active intervention. Extracorporeal shock wave lithotripsy (ESWL) is effective for stones <2 cm, fragmenting calculi for natural passage. Ureteroscopy with laser lithotripsy allows direct visualization and removal of ureteral stones. Percutaneous nephrolithotomy is indicated for large or complex renal stones. Pharmacological therapy may include potassium citrate to alkalinize urine or thiazide diuretics to reduce calcium excretion. Preventive strategies focus on hydration, dietary modification, and treatment of metabolic disorders. Patients are advised to maintain a daily urine output of at least 2–2.5 liters, limit sodium and animal protein intake, and increase dietary citrate. Regular follow-up with imaging and metabolic evaluation helps detect recurrence early. Education on lifestyle modification and adherence to medical therapy is essential to reduce recurrence, prevent renal damage, and improve long-term outcomes.

Conclusion

Urinary stone disease is a prevalent urological disorder that poses significant clinical challenges due to its recurrent nature and potential complications. The etiology is multifactorial, involving metabolic abnormalities, genetic predisposition, dietary factors, dehydration, and urinary tract infections. Clinical manifestations vary, with renal colic, hematuria, and lower urinary tract symptoms being the most common presentations. Accurate diagnosis relies on a combination of laboratory tests and imaging studies, with non-contrast computed tomography recognized as the gold standard. Management depends on stone size, location, and patient factors, ranging from conservative measures and pharmacological therapy to interventional procedures such as extracorporeal shock wave lithotripsy, ureteroscopy, and percutaneous nephrolithotomy. Preventive strategies, including adequate hydration, dietary modifications, and correction of metabolic disturbances, are essential to reduce recurrence and preserve renal function. Early

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

detection, individualized management, and patient education are crucial to improving outcomes, minimizing complications, and enhancing long-term quality of life in individuals affected by urinary stone disease.

References

1. Szymanski, J. R., Chlosta, P., & Przydacz, M. (2025). Epidemiology, diagnostics, and therapeutic strategies in urolithiasis: a comprehensive review. *Przegląd Lekarski*.
2. Jahrreiss, V., Seitz, C., & Quhal, F. (2024). Medical management of urolithiasis: Great efforts and limited progress. *Asian Journal of Urology*, 11(2), 149–155.
3. Razi, A., Ghiaei, A., Kamali Dolatabadi, F., & Haghighi, R. (2024). Unraveling the association of bacteria and urinary stones in patients with urolithiasis: an update review article. *Frontiers in Medicine*.
4. Allam, E. A. H. (2024). Urolithiasis unveiled: pathophysiology, stone dynamics, types, and inhibitory mechanisms: a review. *African Journal of Urology*.
5. Азимова, А. А., Маликов, Д. И., & Шайкулов, Х. Ш. (2021). МОНИТИРОВАНИЕ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ СЕПСИСА ЗА. *PEDAGOGICAL SCIENCES AND TEACHING METHODS*, 48, 18-22.