

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

MANAGEMENT OF SOFT TISSUE INJURIES IN THE MAXILLOFACIAL REGION

Yusufjonov Muxammadkarim

Central Asian Medical University,

Residency in Oral and Maxillofacial Surgery

Abstract: This article examines the clinical features, diagnosis, and treatment principles of soft tissue injuries in the maxillofacial region. The importance of early assessment and proper primary surgical management in preventing complications is emphasized. Particular attention is given to layered wound closure, infection prevention, and preservation of functional structures such as nerves and salivary ducts. Comprehensive treatment includes antibacterial therapy, local wound care, and rehabilitation measures aimed at restoring facial function and appearance. The findings indicate that timely and systematic management significantly reduces infection, scar formation, and functional impairment. Therefore, a combined surgical and rehabilitative approach provides optimal functional and aesthetic outcomes for patients with maxillofacial soft tissue injuries.

Keywords: Maxillofacial injuries, soft tissue trauma, primary surgical treatment, wound management, infection prevention, rehabilitation, facial function.

Introduction

Soft tissue injuries of the maxillofacial region are frequently encountered in oral and maxillofacial surgery due to road accidents, domestic trauma, and sports injuries. Because the facial area is responsible for breathing, mastication, speech, and facial expression, even minor trauma may lead to functional and aesthetic disturbances. The rich blood supply promotes healing but also increases bleeding and edema, while oral microflora raises the risk of infection. Improper management may cause complications such as abscesses, scar deformities, and nerve dysfunction. Therefore, timely diagnosis and appropriate treatment are essential to ensure proper healing and restoration of facial function and appearance.

Relevance

Soft tissue injuries of the maxillofacial region are common in trauma and may cause functional and aesthetic disturbances. Due to rich blood supply and oral microflora, these wounds have a high risk of infection and scarring. Therefore, timely diagnosis and proper treatment remain an important clinical problem.

Main part

Soft tissue injuries of the maxillofacial region present with a wide spectrum of clinical manifestations depending on the depth and mechanism of trauma. The most common findings include pain, bleeding, swelling, and visible tissue disruption. Facial asymmetry may appear when mimetic muscles are affected. Sensory loss indicates possible trigeminal nerve involvement, while facial nerve injury leads to impaired facial expressions. Salivary leakage suggests damage to the parotid duct or minor salivary glands. Limitation of mouth opening may be associated with muscle spasm or temporomandibular joint trauma. Clinical examination begins with evaluation of airway patency and hemodynamic stability. Inspection determines wound size, contamination, and tissue viability. Palpation helps identify foreign bodies and bone fractures. Probing assesses wound depth and communication with the oral cavity. Radiographic imaging is used when bone injury is suspected. Ultrasound assists in evaluation of soft tissues

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

and hematomas. Neurological assessment determines sensory and motor deficits. Laboratory tests help detect inflammatory changes. Early differentiation between simple and complex injuries is essential. Timely diagnosis allows selection of appropriate treatment strategy. Accurate clinical assessment reduces complication risk. Comprehensive examination is therefore fundamental for successful management.

Primary management of maxillofacial soft tissue injuries focuses on prevention of infection and restoration of anatomical integrity. Treatment begins with adequate anesthesia and wound irrigation using sterile saline and antiseptic solutions. Removal of foreign bodies and devitalized tissues is performed carefully to preserve viable structures. Hemostasis is achieved by compression, ligation, or coagulation. Layered closure is essential to restore normal anatomy and function. Muscle repair ensures proper facial movement and symmetry. Special attention is given to repair of salivary ducts and nerves. Skin edges are approximated using atraumatic sutures along natural skin lines. Drainage may be placed in contaminated wounds to prevent hematoma formation. Early closure within the optimal time period improves healing. Antibiotic prophylaxis is commonly administered. Tetanus prophylaxis is also required. Proper surgical technique minimizes scar formation. Adequate wound care instructions are provided to the patient. Follow-up examination evaluates healing progression. Early intervention significantly decreases complication rates. Primary surgical treatment is therefore the key step in management.

Complications of maxillofacial soft tissue injuries may occur if treatment is delayed or inadequate. Infection is the most common early complication and may lead to abscess or cellulitis. Severe cases can spread to deep facial spaces. Scar deformities may develop during late healing stages. Hypertrophic scars and keloids cause aesthetic concerns. Contractures can restrict mouth opening and facial movements. Nerve injuries may result in persistent sensory deficits or facial weakness. Salivary fistulas appear after duct damage. Prevention includes proper surgical technique and antibiotic therapy. Rehabilitation begins soon after wound stabilization. Physiotherapy improves circulation and tissue elasticity. Facial exercises restore muscle function. Massage reduces scar stiffness. Speech therapy helps articulation recovery when necessary. Psychological support improves patient adaptation. Long-term follow-up monitors functional and cosmetic outcomes. Comprehensive rehabilitation enhances quality of life. Successful management depends on both treatment and rehabilitation stages.

Conclusion

Soft tissue injuries of the maxillofacial region require timely diagnosis and a comprehensive treatment approach. Proper primary surgical management with accurate layered repair and infection prevention is essential to ensure optimal healing and to minimize complications. Early intervention significantly reduces the risk of infection, scar deformity, and functional impairment. In addition, rehabilitation measures such as physiotherapy and facial exercises improve recovery of facial function and appearance. Therefore, combined surgical treatment and structured rehabilitation provide the best functional and aesthetic outcomes for patients with maxillofacial soft tissue injuries.

References

1. Alpert, B., & Tiwana, P. (2019). Management of soft tissue injuries in maxillofacial trauma. *Journal of Oral and Maxillofacial Surgery*, 77(5), 1020–1028.
2. Boffano, P., Rocchia, F., Zavattero, E., Dediol, E., Uglešić, V., Kovačić, Z., & European Maxillofacial Trauma Group. (2015). European guidelines for the management of facial trauma. *Journal of Cranio-Maxillofacial Surgery*, 43(6), 935–939.

**Scottish International Conference on Multidisciplinary Research and
Innovation – SICMRI 2026**

3. Chrcanovic, B. R. (2017). Factors influencing infection in facial injuries. *Oral and Maxillofacial Surgery*, 21(1), 23–29.
4. Ellis, E., & Graham, J. (2019). Use of antibiotics in facial fractures and soft tissue injuries. *Journal of Oral and Maxillofacial Surgery*, 77(11), 2300–2306.
5. Fonseca, R. J. (2020). *Oral and maxillofacial trauma* (4th ed.). Elsevier.
6. Kumar, S., & Rahman, S. A. (2020). Primary closure versus delayed closure in contaminated facial wounds. *International Journal of Oral and Maxillofacial Surgery*, 49(9), 1158–1164.