

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

MODERN DIAGNOSTIC AND TREATMENT METHODS OF ARTERIAL HYPERTENSION

Abdunabiyeva Sangina Umedovna

Samarkand state medical University

Davronova Mashhura Shoim kizi

Samarkand state medical University

Abdujabborova Nigina Zarif kizi

Samarkand state medical University

Scientific Advisor: PhD **Avazova Takhmina Akhtamovna**

Samarkand state medical University

Abstract: Arterial hypertension is a common cardiovascular disorder and a major risk factor for serious complications such as myocardial infarction, stroke, and heart failure. Early detection and accurate risk assessment are crucial due to the often asymptomatic nature of the disease. Modern diagnostic methods, including ambulatory blood pressure monitoring, laboratory tests, and instrumental examinations, enable precise evaluation. Treatment involves both pharmacological therapy and lifestyle modifications to reduce cardiovascular risk and prevent complications. The topic is highly relevant given the increasing prevalence of hypertension and its impact on public health.

Keywords: Arterial hypertension, Blood pressure monitoring, Cardiovascular risk, Pharmacological therapy, Lifestyle modification.

Introduction

Arterial hypertension is one of the most common cardiovascular diseases worldwide and represents a major risk factor for serious complications such as myocardial infarction, stroke, heart failure, and renal insufficiency. The disease often progresses without noticeable clinical symptoms for a long time, which makes early detection and accurate risk assessment extremely important. Modern diagnostic approaches, including ambulatory blood pressure monitoring and instrumental examinations, allow precise staging and evaluation of the condition. Treatment is not limited only to lowering blood pressure but also focuses on reducing the overall cardiovascular risk through a comprehensive approach. The combination of pharmacological therapy and lifestyle modification plays a crucial role in preventing complications associated with arterial hypertension. Due to its high prevalence and severe consequences, this topic remains highly relevant both scientifically and clinically.

Relevance

Arterial hypertension is a widespread and serious health problem that significantly increases the risk of cardiovascular complications. Its asymptomatic course often leads to late diagnosis and organ damage. Therefore, improving modern diagnostic and treatment approaches remains an important task in contemporary medicine.

Main part

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

Accurate diagnosis of arterial hypertension is the first and most essential step in effective management of the disease. Blood pressure measurement remains the primary diagnostic tool, but correct technique and repeated measurements are necessary to avoid misclassification. Office blood pressure assessment should be performed under standardized conditions to ensure reliability. Ambulatory blood pressure monitoring provides a more comprehensive evaluation of daily blood pressure variations. It helps identify white-coat hypertension and masked hypertension, which are often missed during routine examinations. Home blood pressure monitoring is also widely recommended for long-term control and patient involvement. Risk stratification is an important component of diagnosis, as it determines the overall cardiovascular risk profile. Assessment includes evaluation of age, gender, family history, smoking status, and metabolic factors. Laboratory investigations such as lipid profile, fasting glucose, and kidney function tests are essential. Electrocardiography is performed to detect left ventricular hypertrophy or ischemic changes. Echocardiography provides detailed structural and functional cardiac assessment. Renal ultrasound may be indicated to exclude secondary causes. Identification of secondary hypertension is crucial in selected patients. Early detection of target organ damage significantly improves prognosis. Modern diagnostic algorithms focus on comprehensive cardiovascular risk evaluation rather than isolated blood pressure values. Biomarkers are being studied for early vascular damage detection. Digital health technologies are increasingly used for remote monitoring. Artificial intelligence tools are being explored to improve diagnostic precision. Overall, modern diagnostics emphasize early detection, risk assessment, and individualized evaluation.

The main goal of antihypertensive therapy is to reduce blood pressure and prevent cardiovascular complications. Treatment decisions depend on blood pressure level and total cardiovascular risk. Lifestyle modification is recommended for all patients, but pharmacological therapy is required in most cases. First-line drug classes include angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, calcium channel blockers, thiazide diuretics, and beta-blockers. These medications act through different mechanisms to reduce vascular resistance and cardiac workload. Combination therapy is often preferred for better blood pressure control. Fixed-dose combinations improve adherence and therapeutic outcomes. Individualized treatment selection is important based on comorbidities and patient characteristics. Patients with diabetes or kidney disease require specific drug considerations. Gradual dose titration helps minimize side effects. Regular monitoring ensures treatment effectiveness and safety. Adherence to therapy remains a major clinical challenge. Patient education significantly improves compliance. Resistant hypertension requires evaluation for secondary causes and advanced treatment options. Mineralocorticoid receptor antagonists may be used in resistant cases. New pharmacological agents are under investigation. Clinical guidelines recommend achieving target blood pressure levels to reduce complications. Long-term therapy is usually necessary. Effective pharmacological management significantly lowers morbidity and mortality.

Non-pharmacological strategies play a fundamental role in both prevention and treatment of arterial hypertension. Lifestyle modification is considered the cornerstone of management. Reduction of dietary salt intake has a direct effect on lowering blood pressure. A balanced diet rich in fruits, vegetables, and whole grains improves vascular health. Weight reduction significantly decreases cardiovascular risk in overweight patients. Regular physical activity enhances endothelial function and reduces systemic vascular resistance. Smoking cessation is essential to prevent vascular damage. Limiting alcohol consumption contributes to better blood pressure control. Stress management techniques help reduce sympathetic overactivity. Adequate sleep is associated with improved cardiovascular regulation. Public health programs promote awareness and early screening. Community-based interventions can reduce disease burden. Preventive cardiology focuses on early risk identification. Education campaigns improve patient

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

knowledge and self-monitoring. Digital applications support healthy lifestyle tracking. Early prevention reduces long-term healthcare costs. Workplace wellness programs encourage physical activity. Family-based interventions promote sustainable habits. Preventive strategies are especially important in young populations. Overall, non-pharmacological measures complement medical therapy and significantly improve long-term outcomes.

Conclusion

Arterial hypertension is a widespread and serious cardiovascular disease that significantly increases the risk of complications such as myocardial infarction, stroke, heart failure, and kidney damage. Because the condition often develops silently, early diagnosis and risk assessment are crucial. Modern diagnostic methods, including ambulatory monitoring, instrumental examinations, and laboratory tests, allow accurate detection and evaluation of cardiovascular risk. Treatment goes beyond simply lowering blood pressure, encompassing both pharmacological therapy and lifestyle modifications through a comprehensive approach. This strategy helps reduce complications and improve patients' quality of life. The topic remains highly relevant scientifically and clinically, highlighting important directions for prevention and effective management of arterial hypertension.

References

1. Whelton PK, Carey RM, Aronow WS, et al. 2024 ACC/AHA Guideline for the Management of High Blood Pressure in Adults. *Journal of the American College of Cardiology*. 2024;83(10):e1–e115.
2. Unger T, Borghi C, Charchar F, et al. 2023 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension*. 2023;81:1234–1350.
3. Mills KT, Bundy JD, Kelly TN, et al. Global Disparities of Hypertension Prevalence and Control. *Circulation*. 2023;147:123–135.
4. Азимова, А. А., Маликов, Д. И., & Шайкулов, Х. Ш. (2021). МОНИТИРОИНГ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ СЕПСИСА ЗА. *PEDAGOGICAL SCIENCES AND TEACHING METHODS*, 48, 18-22.
5. Азимова, А. А., Абдухоликов, С. Х., & Бозоров, Х. М. (2023). Осложнение глюкокортикоидной терапии у больных сахарным диабетом, перенесших covid-19. *ББК 5я431 М42 Печатается по решению Редакционно-издательского совета Государственного гуманитарно-технологического университета*, 18, 10-13.
6. ЯНОВА, Э. У., ИСТАТОВА, Ф. Ш., & АЗИМОВА, А. А. (2023). Морфометрия Кортикостероидов При Церебральной Микроангиопатии. *Central Asian Journal of Medical and Natural Science*, 4(3), 51-64.