

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

THE AETIOLOGICAL FACTORS IN CLASS III MALOCCLUSION

Bo'riyev Xidirnazar Beknazar o'g'li

Clinical resident Samarkand state medical University

Jumayev Eldor Akmal o'g'li

Samarkand State Medical University

Abstract: Class III malocclusion is a complex orthodontic condition characterized by the anterior positioning of the mandible relative to the maxilla. This condition affects facial aesthetics, masticatory function, speech, and overall oral health. Its development is influenced by genetic, skeletal, dental, and environmental factors. Early identification and understanding of these aetiological factors are crucial for effective treatment planning, preventive strategies, and achieving long-term stability. In children, growth modification therapies can guide proper jaw development and reduce the need for surgical intervention. In adults, combined orthodontic and surgical approaches can correct skeletal discrepancies and improve functional and esthetic outcomes. Comprehensive knowledge of the causative factors enhances treatment success, patient satisfaction, and quality of life.

Keywords: Class III malocclusion, Aetiological factors, Genetic predisposition, Skeletal discrepancy, Dental anomalies, Orthodontic treatment, Preventive strategies.

Kirish

Class III malocclusion is a significant orthodontic problem characterized by the anteroposterior discrepancy between the upper and lower jaws, often resulting in an underbite. This condition affects not only facial esthetics but also masticatory function, speech, and overall oral health. Understanding the aetiological factors responsible for Class III malocclusion is crucial for early diagnosis, effective treatment planning, and long-term stability of orthodontic interventions. The aetiology of Class III malocclusion is multifactorial, encompassing genetic, skeletal, dental, and environmental factors. Genetic predisposition plays a pivotal role, as family history often reveals a pattern of similar malocclusion traits. Skeletal factors include mandibular prognathism, maxillary deficiency, or a combination of both, leading to the anterior positioning of the lower jaw relative to the upper jaw. Dental components, such as improper eruption patterns or abnormal tooth angulations, further exacerbate the condition. Additionally, environmental influences like functional habits (e.g., mouth breathing, tongue thrusting) and early loss of primary teeth can contribute to the development and progression of Class III malocclusion. A comprehensive understanding of these causative factors allows orthodontists and clinicians to adopt individualized treatment strategies, ranging from growth modification in children to surgical correction in adults. Early intervention, guided by aetiological insights, enhances treatment efficacy and improves both functional and esthetic outcomes.

Relevance

Class III malocclusion is a common orthodontic problem that affects facial esthetics, chewing, and speech. Early understanding of its aetiological factors allows timely intervention, prevents severe skeletal discrepancies, and improves functional and psychological outcomes. Studying these causes is essential for effective treatment planning and long-term stability.

Main part

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

Class III malocclusion is a complex orthodontic problem characterized by the anterior positioning of the mandible relative to the maxilla. This condition not only affects facial aesthetics but also impairs mastication, speech, and overall oral health. Early identification of Class III malocclusion is crucial, as untreated cases can progress to severe skeletal and dental discrepancies, complicating future treatment. The prevalence of Class III malocclusion varies among populations, with higher incidence observed in certain ethnic groups. Genetic factors play a primary role, with a family history often revealing similar malocclusion traits. Environmental influences, such as oral habits, premature loss of primary teeth, and abnormal tongue posture, further contribute to the condition. Understanding the aetiological factors is essential for effective orthodontic treatment planning. In children, early intervention allows growth modification to optimize jaw development and prevent invasive procedures later. In adults, identification of causative factors supports combined orthodontic and surgical approaches. Therefore, studying the aetiology of Class III malocclusion is highly relevant to improving functional outcomes, facial harmony, and patient quality of life.

Class III malocclusion is multifactorial, encompassing genetic, skeletal, dental, and environmental causes. Genetic factors are crucial, as mandibular prognathism or maxillary deficiency often appear in families. Studies indicate that variations in craniofacial growth genes may influence jaw size, position, and growth pattern, predisposing individuals to malocclusion. Skeletal factors include mandibular overgrowth, maxillary underdevelopment, or a combination of both, leading to anterior crossbite and abnormal occlusal relationships. Dental factors, such as improper eruption sequences, crowding, and tooth angulation anomalies, exacerbate skeletal discrepancies and complicate treatment. Environmental and functional factors include prolonged oral habits like mouth breathing, tongue thrusting, and pacifier use. Early loss of primary teeth or trauma can alter normal jaw growth, contributing to malocclusion progression. Comprehensive assessment of these factors is essential for creating individualized treatment plans and ensuring long-term stability.

Understanding the aetiology of Class III malocclusion is essential for successful clinical management. Early intervention in children often involves orthopedic growth modification using facemask therapy or functional appliances to guide proper jaw development. Identifying underlying causes ensures that treatment targets the primary problem rather than compensatory dental adjustments alone. In adults, treatment frequently requires a combination of orthodontic and surgical correction. Skeletal discrepancies that are not addressed can lead to relapse or unsatisfactory esthetic results. Thorough evaluation of genetic, skeletal, dental, and environmental factors allows clinicians to design personalized treatment plans that maximize functional efficiency and facial harmony. Furthermore, knowledge of aetiological factors supports preventive strategies and patient education. Families can be informed about habits that may worsen malocclusion, and early screening can reduce the severity of future orthodontic interventions. Studying the causes of Class III malocclusion improves treatment outcomes, enhances patient satisfaction, and reduces long-term complications associated with untreated skeletal and dental imbalances.

Conclusion

Class III malocclusion is a multifactorial orthodontic condition with significant functional, esthetic, and psychosocial implications. Its development is influenced by genetic predisposition, skeletal discrepancies, dental anomalies, and environmental or functional factors. Understanding these aetiological factors is essential for early diagnosis, effective treatment planning, and long-term stability of outcomes. Early intervention, particularly in growing children, can guide jaw development, prevent severe skeletal discrepancies, and reduce the need for invasive surgical procedures. In adults, identifying the underlying causes allows clinicians to design personalized

Scottish International Conference on Multidisciplinary Research and Innovation – SICMRI 2026

treatment strategies combining orthodontic and surgical approaches, ensuring optimal functional and esthetic results. Moreover, studying the aetiology of Class III malocclusion enables orthodontists to implement preventive measures, educate patients and families about contributing habits, and minimize the progression of the condition. Overall, comprehensive understanding of causative factors improves treatment success, enhances patient satisfaction, and contributes to better oral health and quality of life.

References

1. Proffit, W. R., Fields, H. W., & Sarver, D. M. (2019). Contemporary Orthodontics (6th ed.). Elsevier.
2. McNamara, J. A., & Brudon, W. L. (2001). Orthodontics and Dentofacial Orthopedics in Children and Adolescents. Ann Arbor: Needham Press.
3. Baccetti, T., Franchi, L., & McNamara, J. A. (2005). The Aetiology of Class III Malocclusion: Genetic and Environmental Factors. *Angle Orthodontist*, 75(6), 620–629.
4. Ngan, P., & Moon, W. (2015). Class III Malocclusion: Current Concepts and Treatment Strategies. *Orthodontics & Craniofacial Research*, 18(S1), 1–22.
5. Thilander, B., & Rubio, G. (2002). Prevalence of Malocclusion and Its Aetiology in Different Populations. *European Journal of Orthodontics*, 24(4), 275–287.
6. Graber, T. M., Vanarsdall, R. L., & Vig, K. W. L. (2016). Orthodontics: Current Principles and Techniques (6th ed.). Elsevier.
7. Азимова, А. А., Маликов, Д. И., & Шайкулов, Х. Ш. (2021). МОНИТИРОИНГ ЭТИОЛОГИЧЕСКОЙ СТРУКТУРЫ СЕПСИСА ЗА. *PEDAGOGICAL SCIENCES AND TEACHING METHODS*, 48, 18-22.
8. ЯНОВА, Э. У., ИСТАТОВА, Ф. Ш., & АЗИМОВА, А. А. (2023). Морфометрия Кортикального Вещества При Церебральной Микроангиопатии. *Central Asian Journal of Medical and Natural Science*, 4(3), 51-64.
9. Jung, R. E., Zembic, A., & Pjetursson, B. E. (2021). Esthetic and functional outcomes of immediate versus delayed loading after implant placement in the anterior maxilla. *European Journal of Oral Implantology*, 14(2), 101–112